

Chemistry 11

Solution Practice Test

Name: *Key*
 Date:
 Block:

1. During a lab activity, you dilute 45 mL of a 8.5 M HCl solution to a final volume of 120 mL. What is the resulting HCl concentration?

- a. 2.3 M
- b. 3.2 M
- c. 5.1 M
- d. 23 M

$$C_1V_1 = C_2V_2$$

$$(8.5)(45) = C_2(120)$$

$$C_2 = 3.2M$$

2. Which of the following equations correctly represents the dissociation of calcium chloride in water?

- a. $\text{CaCl}_{(aq)} \rightarrow \text{Ca}^{+(aq)} + \text{Cl}^{-(aq)}$
- b. $\text{CaCl}_{2(aq)} \rightarrow \text{Ca}^{2+(aq)} + \text{Cl}^{-(aq)}$
- c. $\text{CaCl}_{2(aq)} \rightarrow \text{Ca}^{2+(aq)} + 2\text{Cl}^{-(aq)}$
- d. $\text{CaCl}_{2(aq)} + \text{H}_2\text{O} \rightarrow \text{Ca}^{2+(aq)} + 2\text{Cl}^{-(aq)}$



3. When added to a solution containing Mg^{2+} , which anion will create a precipitate?

- a. NO_3^-
- b. Cl^-
- c. OH^-
- d. S^{2-}

4. When added to an iron (III) iodide solution, which of the following compounds will create a precipitate?

- a. cesium nitrate
- b. hydrochloric acid
- c. copper (II) sulphate
- d. ammonium hydroxide



5. How many mL of 0.550 M NaOH would be required to titrate 25.0 mL of a 0.388 M solution of hydrochloric acid?

- a. 17.6 mL
- b. 25.0 mL
- c. 35.4 mL
- d. 46.9 mL



$$25.0 \text{ mL HCl} \times \frac{1 \text{ L}}{1000 \text{ mL}} \times \frac{0.388 \text{ mol HCl}}{1 \text{ L}} \times \frac{1 \text{ mol NaOH}}{1 \text{ mol HCl}} \times \frac{1 \text{ L}}{0.550 \text{ mol NaOH}} \times \frac{1000 \text{ mL}}{1 \text{ L}} = 17.6 \text{ mL NaOH}$$

6. A student must prepare a 3.00 L solution of 0.750 M NaOH. She is given a stock solution of 12.0 M NaOH.

a. What volume of stock solution is needed to prepare the final solution?

$$C_1V_1 = C_2V_2$$

$$(3.00)(0.750) = (12.0)V_2$$

$$V_2 = \boxed{0.188 \text{ L NaOH}}$$



b. What volume of water was added to prepare the final solution?

$$3.00 \text{ L} - 0.188 \text{ L} = \boxed{2.812 \text{ H}_2\text{O}}$$

7. MnSO_4 was dissolved in water. What is the ionization equation?



8. A chemist mixes a 225 mL of a 3.8 M Na_2CO_3 solution with 3.8 g of K_2CO_3 . What is the concentration of each ion in this solution?



$$[\text{Na}^+] = 7.6 \text{ M}$$

$$[\text{CO}_3^{2-}] = 3.8 \text{ M} + 0.12 \text{ M} = 3.9 \text{ M}$$

$$[\text{K}^+] = 0.24 \text{ M}$$

$$[\text{K}_2\text{CO}_3] \frac{3.8 \text{ g K}_2\text{CO}_3}{225 \text{ mL}} \times \frac{1000 \text{ mL}}{1 \text{ L}} \times \frac{1 \text{ mol K}_2\text{CO}_3}{138.21 \text{ g}}$$


$$= 0.12 \text{ M K}_2\text{CO}_3$$

$$\text{K}_2\text{CO}_3 \rightarrow 2\text{K}^+ + \text{CO}_3^{2-}$$


0.12 M 0.24 M 0.12 M

9. For the following solutions, use a flow chart to describe the process of separating the ions from each other.



a. Mg^{2+} , Pb^{2+} and Zn^{2+}



b. OH^- , PO_4^{3-} , S^{2-}




10. Predict the products in the following reactions. Then, balance the equations. Be sure to indicate the state (aq or s), of each product.




11. Write a formula equation, complete ionic equation and net ionic equation for the following reactions:

a. Strontium hydroxide and zinc chloride



b. Ammonium bromide and copper (I) sulphate



12. Determine whether the following are soluble or have low solubility.

a. Barium chloride

soluble / low solubility (circle one)

b. $\text{Sn}(\text{OH})_4$

soluble / low solubility (circle one)

13. A titration was performed that required 12.7 mL of 0.150 M $\text{Mg}(\text{OH})_2$ to titrate 25.00 mL of a hydrochloric acid, HCl, solution. Determine the molarity of the hydrochloric acid.



$$12.7 \text{ mL} \times \frac{1 \text{ L}}{1000 \text{ mL}} \times \frac{0.150 \text{ mol Mg}(\text{OH})_2}{1 \text{ L}} \times \frac{2 \text{ mol HCl}}{1 \text{ mol Mg}(\text{OH})_2}$$

$$= 0.00381 \text{ mol HCl}$$

$$\frac{0.00381 \text{ mol HCl}}{25.00 \text{ mL}} \times \frac{1000 \text{ mL}}{1 \text{ L}} = \boxed{0.152 \text{ M HCl}}$$

14. Consider the following results from a titration lab.

4.50 g of KOH was dissolved to make a 100. mL solution
Below is the volume of the KOH solution needed to neutralize 15.0 mL H₃PO₄.

	Trial #1	Trial #2	Trial #3	Trial #4
Initial reading of burette (mL)	2.56	8.95	13.35	17.55
Final reading of burette (mL)	8.95	13.35	17.55	21.75
Volume of KOH (mL)	6.39 mL	4.40 mL	4.20 mL	4.20 mL

a. What is the concentration of the standardized solution of KOH?

$$\frac{4.50 \text{ g KOH}}{100. \text{ mL}} \times \frac{1000 \text{ mL}}{1 \text{ L}} \times \frac{1 \text{ mol KOH}}{56.11 \text{ g}} = \boxed{0.802 \text{ M KOH}}$$

b. What was the average volume of KOH was needed? (Only use data from three trials!!)

** use closest 3 trials!!*

$$\frac{4.40 \text{ mL} + 4.20 \text{ mL} + 4.20 \text{ mL}}{3} = \boxed{4.27 \text{ mL KOH}}$$

c. What is the concentration of the acid?



$$4.27 \text{ mL KOH} \times \frac{1 \text{ L}}{1000 \text{ mL}} \times \frac{0.802 \text{ mol KOH}}{1 \text{ L}} \times \frac{1 \text{ mol H}_3\text{PO}_4}{3 \text{ mol KOH}} = 0.00114 \text{ mol H}_3\text{PO}_4$$

$$\frac{0.00114 \text{ mol H}_3\text{PO}_4}{15.0 \text{ mL}} \times \frac{1000 \text{ mL}}{1 \text{ L}} = \boxed{0.0760 \text{ M H}_3\text{PO}_4}$$